

1. GENERAL INFORMATION

IUPAC Name: 3-(2-ethylphenyl)-2-methylquinazolin-4(3H)-one

CFR: Not Scheduled (9/2013)

CAS#: 7432-25-9

Synonyms: Aolan, Athinazone, Ethinazone, NSC 96163

Source: DEA Reference Material Collection

Appearance: White powder (HCl)

Kovat's Index: Pending

UV_{max} (nm): 226.1, 265.3

2. CHEMICAL AND PHYSICAL DATA

2.1 CHEMICAL DATA

Form	Chemical Formula	Molecular Weight	Melting Point (°C)
Base	C ₁₇ H ₁₆ N ₂ O	264	Not Determined
HCl	C ₁₇ H ₁₆ N ₂ O · HCl	300	265.1

3. ADDITIONAL RESOURCES

[Forendex](#)

[Wikipedia](#)

4. QUALITATIVE DATA

4.1 NUCLEAR MAGNETIC RESONANCE

Method NMR CD₃OD

Sample Preparation: Dilute analyte to ~10 mg/mL in CD₃OD containing TMS for 0 ppm reference and dimethylsulfone as quantitative internal standard.


Instrument: 400 MHz NMR spectrometer

Parameters: Spectral width: at least containing -3 ppm through 13 ppm

Pulse angle: 90°

Delay between pulses: 45 seconds

¹H NMR: Etaqualone HCl; Lot N17-P55C; CD₃OD; 400 MHz



4.2 Gas Chromatography/Mass Spectrometry

Sample Preparation: Dilute analyte ~ 1 mg/mL in methanol

Instrument: Agilent gas chromatograph operated in split mode with MS detector

Column: DB-1 MS (or equivalent); 30m x 0.25 mm x 0.25 μ m

Carrier Gas: Helium at 1 mL/min

Temperatures: Injector: 280°C

MSD transfer line: 280°C

MS Source: 230°C

MS Quad: 150°C

Oven program:

1) 100°C initial temperature for 1.0 min

2) Ramp to 300°C at 12 °C/min

3) Hold final temperature for 9.0 min

Injection Parameters: Split Ratio = 20:1, 1 μ L injected

MS Parameters: Mass scan range: 30-550 amu


Threshold: 100

Tune file: stune.u

Acquisition mode: scan

Retention Time: 13.853 min

EI Mass Spectrum: Etaqualone HCl; Lot 5667-1022-74



4.3 INFRARED SPECTROSCOPY (FTIR)

Instrument: FTIR with diamond ATR attachment (3 bounce)

Scan Parameters: Number of scans: 32
Number of background scans: 32
Resolution: 4 cm⁻¹
Sample gain: 8
Aperture: 150

FTIR ATR (Diamond, 3 Bounce): Etaqualone HCl; Lot N17-P55C

